



**LEVOTHYROXINE USE IN THE ELDERLY IS ASSOCIATED WITH AN INCREASED RISK OF FRACTURES .....2**

Turner MR, Camacho X, Fischer HD, Austin PC, Anderson GM, Rochon RA, Lipscombe LL. **Levothyroxine dose and risk of fractures in older adults: nested case-control study.** BMJ 2011;342:d2238.

**SUBCLINICAL HYPOTHYROIDISM IS ASSOCIATED WITH INCREASED CORONARY HEART DISEASE AND ALL-CAUSE MORTALITY. ....4**

McQuade C, Skugor M, Brennan DM, Hoar B, Stevenson C, Hoogwerf BJ. **Hypothyroidism and Moderate Subclinical Hypothyroidism Are Associated with Increased All-Cause Mortality Independent of Coronary Heart Disease Risk Factors: A PreCIS Database Study.** Thyroid 2011;21:837-43. Epub July 11, 2011.

**WOMEN OF CHILDBEARING AGE MAY BENEFIT FROM IODINE SUPPLEMENTATION SEVERAL MONTHS BEFORE CONCEPTION .....6**

Moleti M, Di Bella B, Giorgianni G, Mancuso A, De Vivo A, Alibrandi A, Trimarchi F, Vermiglio F. **Maternal thyroid function in different conditions of iodine nutrition in pregnant women exposed to mild-moderate iodine deficiency: an observational study.** Clin Endocrinol (Oxf) 2011;74:762-8.

**TREATMENT OF GOITER AND NODULES IS SUCCESSFUL WITH NON-TSH-SUPPRESSING DOSES OF THYROXINE PLUS IODINE .....9**

Grussendorf M, Reiners C, Paschke R, Wegscheider K, on behalf of the LISA investigators. **Reduction of thyroid nodule volume by levothyroxine and iodine alone and in combination: a randomized, placebo-controlled trial.** J Clin Endocrinol Metab. June 29, 2011 [Epub ahead of print]. doi: 10.1210/jc.2011-0356.

**SMALL THYROID BED MASSES FOUND AFTER INITIAL TREATMENT OF DIFFERENTIATED THYROID CANCER HAVE A BENIGN OUTCOME. .... 11**

Rondeau G, Fish S, Hann LE, Fagin JA, Tuttle RM. **Ultrasonographically detected small thyroid bed nodules identified after total thyroidectomy**

**for differentiated thyroid cancer seldom show clinically significant structural progression.**

Thyroid 2011;21:845-53.

**ARE LOBECTOMY AND THYROIDECTOMY WITHOUT RADIOACTIVE IODINE REMNANT ABLATION A SAFE PROCEDURE IN SELECTED THYROID CANCER PATIENTS? ..... 13**

Vaisman F, Shaha A, Fish S, Tuttle R. **Initial therapy with either thyroid lobectomy or total thyroidectomy without radioactive iodine remnant ablation is associated with very low rates of structural disease recurrence in properly selected patients with differentiated thyroid cancer.** Clin Endocrinol (Oxf). February 8, 2011 [Epub ahead of print]. doi: 10.1111/j.1365-2265.2011.04002.x.

**THE DOUBLING TIME OF SERUM THYROGLOBULIN IS A VERY STRONG PREDICTOR OF PROGNOSIS IN PATIENTS WITH PAPILLARY THYROID CARCINOMA. .... 15**

Miyauchi A, Kudo T, Miya A, Kobayashi K, Ito Y, Takamura Y, Higashiyama T, Fukushima M, Kihara M, Inoue H, Tomoda C, Yabuta T, Masuoka H. **Prognostic impact of serum thyroglobulin doubling-time under thyrotropin suppression in patients with papillary thyroid carcinoma who underwent total thyroidectomy.** Thyroid 2011;21:707-16. Epub June 7, 2011.

**MEDULLARY THYROID MICROCARCINOMAS HAVE SIGNIFICANT RATES OF POOR PROGNOSTIC FEATURES AND REQUIRE APPROPRIATE SURGICAL MANAGEMENT. .... 17**

Kazaure HS, Roman SA, Sosa JA. **Medullary thyroid microcarcinoma: a population-level analysis of 310 patients.** Cancer. June 29, 2011 [Epub ahead of print]. doi: 10.1002/cncr.26283.

**WHEN IS A POSITIVE ANTI-TG, ANTI-TPO OR ANTI-TSH-RECEPTOR TITER CLINICALLY USEFUL? ..... 19**

Hutfless S, Matos P, Talor MV, Caturegli P, Rose NR. **Significance of prediagnostic thyroid antibodies in women with autoimmune thyroid disease.** J Clin Endocrinol Metab. June 29, 2011 [Epub ahead of print]. doi:10.1210/jc.2011-0228.










## SUBCLINICAL HYPOTHYROIDISM IS ASSOCIATED WITH INCREASED CORONARY HEART DISEASE AND ALL-CAUSE MORTALITY


### References

1. Imaizumi M, Akahoshi M, Ichimaru S, Nakashima E, Hida A, Soda M, Usa T, Ashizawa K, Yokoyama N, Maeda R, et al. Risk for ischemic heart disease and all-cause mortality in subclinical hypothyroidism. *J Clin Endocrinol Metab* 2004;89:3365-70.
2. Walsh JP, Bremner AP, Bulsara MK, O’Leary P, Leedman PJ, Feddema P, Michelangeli V. Subclinical thyroid dysfunction as a risk factor for cardiovascular disease. *Arch Intern Med* 2005;165:2467-72.
3. Iervasi G, Molinaro S, Landi P, Taddei MC, Galli E, Mariani F, L’Abbate A, Pingitore A. Association between increased mortality and mild thyroid dysfunction in cardiac patients. *Arch Intern Med* 2007;167:1526-32.


DEDICATED TO SCIENTIFIC INQUIRY, CLINICAL EXCELLENCE, PUBLIC SERVICE, EDUCATION, AND COLLABORATION.




AMERICAN THYROID ASSOCIATION  
FOUNDED 1923



ATA Publications



Public & Patients



Physicians & Professionals

[www.thyroid.org](http://www.thyroid.org)

ABOUT THE ATA    GIVE ONLINE    JOIN THE ATA    FELLOWS' CORNER    MEMBERS ONLY

## We invite you to join the ATA!

### Are You Intrigued by the Study of the Thyroid? You Belong in the ATA!

- ATA members are leaders in thyroidology who promote excellence and innovation in clinical care, research, education, and public policy.
- Join us as we advance our understanding of the causes and improve the clinical management of thyroid diseases in this era of rapid pace biomedical discovery.
- A close-knit, collegial group of physicians and scientists, the ATA is dedicated to the research and treatment of thyroid diseases. ATA's rich history dates back to 1923 and its members are respected worldwide as leaders in thyroidology.
- The ATA encourages you to apply for membership. We want you to experience the wealth of knowledge and enjoy the benefits of being active in this highly specialized and regarded society. The ATA looks forward to having you as a member!





## WOMEN OF CHILDBEARING AGE MAY BENEFIT FROM IODINE SUPPLEMENTATION SEVERAL MONTHS BEFORE CONCEPTION

incidence of hypothyroidism in pregnancy is estimated to be around 4%, with most of the women having subclinical disease; the women who were detected were newly diagnosed or those with iodine deficiency (the incidence is very low in the United States), or about 40% of women who were on levothyroxine therapy at the time of the first obstetrical visit (4). It was shown that women on levothyroxine (L-T<sub>4</sub>) therapy with a preconception serum TSH <1.3 mIU/L (5), attained a normal serum TSH (<2.5 mIU/L) at the first obstetrical visit. Therefore, it appears reasonable, until further studies confirm the work of Moleti et al.,

to advise all women in the United States who are of reproductive age to add an extra 150 µg of iodine daily to their regular diet, and in addition to advise those on L-T<sub>4</sub> therapy to maintain their serum TSH levels at not more than 1.3 mIU/L. The exception is women who have undergone thyroidectomy for thyroid cancer, who usually require a lower serum TSH level. As we all know very well, unplanned pregnancy is not a rare event in our daily practice.

— Jorge H. Mestman, MD

### References

1. Zimmermann MB. The impact of iodised salt or iodine supplements on iodine status during pregnancy, lactation and infancy. *Public Health Nutr* 2007;10:1584-5.
2. Glinioer D. The regulation of thyroid function in pregnancy: pathways of endocrine adaptation from physiology to pathology. *Endocr Rev* 1997;18:404-31.
3. Pierce E. Commentary on Caldwell KL, Makhmudov A, Ely E, Jones RL, Wang RY. Iodine status of the U.S. population, National Health and Nutrition Examination Survey, 2005–2006 and 2007–2008. *Clin Thyroidol* 2011;23:2-3.
4. McClain MR, Lambert-Messerlian G, Haddow JE, Palomaki GE, Canick JA, Cleary-Goldman J, Malone FD, Flint Porter TF, Nyberg DA, Bernstein P, et al. Sequential first- and second-trimester TSH, free thyroxine, and thyroid antibody measurements in women with known hypothyroidism: a FaSTER trial study. *Am J Obstet Gynecol* 2008;129:e1-6. Epub April 29, 2008.
5. Abalovich M, Alcaraz G, Kleiman-Rubinsztein J, Pavlove MM, Cornelio C, Levalle O, Gutierrez S. The relationship of preconception thyrotropin levels to requirements for increasing the levothyroxine dose during pregnancy in women with primary hypothyroidism. *Thyroid* 2010;20:1175-8.

















## THE DOUBLING TIME OF SERUM THYROGLOBULIN IS A VERY STRONG PREDICTOR OF PROGNOSIS IN PATIENTS WITH PAPILLARY THYROID CARCINOMA

the first four data points was a significant prognostic factor in univariate and multivariate analysis. In this study, thyroid cancer-specific deaths occurred only in stage IV disease. But only 6 of 189 (3.2%) of patients in stage IV died of the disease, giving a 10-year cause-specific survival rate of 94.6%. Using the Tg-DT, 5 of 17 patients with a Tg-DT of <1 year and 1 of 21 of patients with Tg-DT of 1 to 3 years died of thyroid cancer, while none of the patients in the other groups died. Thus, the Tg-DT was better than TMN staging at

predicting the risk of death. This investigation has put a quantitative number on what we are already knew clinically, namely that patients with rising Tg levels are at high risk for recurrence and death. We should use the Tg-DT in the same way we use calcitonin doubling time to predict which patients with medullary thyroid cancer are at high risk for recurrence and death.

— **Stephanie L. Lee, MD, PhD**

### References

1. Baudin E, Do Cao C, Cailleux AF, Leboulleux S, Travagli JP, Schlumberger M. Positive predictive value of serum thyroglobulin levels, measured during the first year of follow-up after thyroid hormone withdrawal, in thyroid cancer patients. *J Clin Endocrinol Metab* 2003;88:1107-11.
2. Miyauchi A, Onishi T, Morimoto S, Takai S, Matsuzuka F, Kuma K, Maeda M, Kumahara Y. Relation of doubling time of plasma calcitonin levels to prognosis and recurrence of medullary thyroid carcinoma. *Ann Surg* 1984;199:461-6.



American Thyroid Association

Prevent  
Diagnose  
Treat

[www.thyroid.org](http://www.thyroid.org)

Support valuable patient education  
and crucial thyroid research!





## MEDULLARY THYROID MICROCARCINOMAS HAVE SIGNIFICANT RATES OF POOR PROGNOSTIC FEATURES AND REQUIRE APPROPRIATE SURGICAL MANAGEMENT

The study raises the question of screening small nodules by measuring serum calcitonin in order to detect MTC at an early stage. This is a debatable topic because elevated serum calcitonin detected a 0.5 to 1.5% incidence of microMTC in several large European series of patients who were going to have surgery for nodular goiter, but not for small nodules. The data are summarized well in an editorial by Hodak and Burman, who concluded that calcitonin screening without evidence of a family history of

MTC yielded too many false positives associated with thyroiditis (1). In a review by Valle and Kloos of 24 autopsy series published from 21 countries, the average prevalence of occult microMTC was 0.14% (2). Finally, the current ATA guidelines “cannot recommend either for or against the routine measurement of serum calcitonin” for evaluation of thyroid nodules (3).

— Jerome M. Hershman, MD

### References

1. Hodak SP, Burman KD. Editorial: The calcitonin conundrum—is it time for routine measurement of serum calcitonin in patients with thyroid nodules? *J Clin Endocrinol Metab* 2004;89:511-4.
2. Valle LA, Kloos RT. The prevalence of occult medullary thyroid carcinoma at autopsy. *J Clin Endocrinol Metab*. 2011;96:E109-E113.
3. Cooper DS, Doherty GM, Haugen BR, Kloos RT, Lee SL, Mandel SJ, Mazzaferri EL, McIver B, Pacini F, Schlumberger M, et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid*. 2009;19:1167-214.



# WHEN IS A POSITIVE ANTI-TG, ANTI-TPO OR ANTI-TSH-RECEPTOR TITER CLINICALLY USEFUL?

Hashimoto’s thyroiditis often presents as a goiter in adolescents and young adults, and these patients often remain euthyroid, whereas some whose TSH was initially above normal or who were overtly hypothyroid subsequently return to euthyroidism, and yet others continue to or subsequently have hypothyroidism (1). Adult patients with Hashimoto’s thyroiditis often do not have a goiter (clinical information about the patients was not provided in the present study). A common diagnostic and clinical problem is the patient who has symptoms that might indicate hypothyroidism but who has no positive physical findings. When tests for T<sub>4</sub>/free T<sub>4</sub>, T<sub>3</sub>, and TSH are ordered, they turn out to be normal. However, if antithyroid antibodies are also ordered and they turn out to be positive, what should the physician do? Most will follow the patient’s TSH level closely. However, if you simply assume that the diagnosis is Hashimoto’s thyroiditis and put the patient on levothyroxine therapy, then even if the patient’s symptoms improve, you have not established that your diagnosis is correct. On the other hand, a euthyroid woman with positive antibodies and a TSH that is within the normal range

but >2.5 mU/L is more than four times as likely to have hypothyroidism over the next 13 years than if her TSH is ≤2.5 mU/L (2).

Only 55% of the patients with Graves’ disease had positive anti-TSH-R antibodies in this study, perhaps because samples could be drawn 6 months after the diagnosis was made, and antithyroid therapy can reduce mean anti-TSHR-stimulating antibody levels by more than half (and anti-TPO titers by two-thirds) in adults within 6 months (3). Anti-TSH-R assays can be useful in evaluating pregnant women who currently have or previously had Graves’ disease; in newborns with possible neonatal hyperthyroidism or who may be transiently hypothyroid because of blocking antibodies; in patients in whom euthyroid Graves’ ophthalmopathy is suspected; in confirming Graves’ disease in hyperthyroid patients in whom a radioiodine uptake and scan should not or cannot be performed; and possibly in determining the likelihood of a recurrence of Graves’ disease before discontinuing antithyroid drug therapy.

— Stephen W. Spaulding, MD

### References

1. Demirbilek H, Kandemir N, Gonc EN, et al. Assessment of thyroid function during the long course of Hashimoto’s thyroiditis in children and adolescents. *Clin Endocrinol* 2009;71:451-4.
2. Walsh JP, Bremner AP, Feddema P, et al. Thyrotropin and thyroid antibodies as predictors of hypothyroidism: a 13-year, longitudinal study of a community-based cohort using current immunoassay techniques. *J Clin Endocrinol Metab* 2010;95:1095-104.
3. Guilhem I, Massart C, Poirier JY, Maugendre D. Differential evolution of thyroid peroxidase and thyrotropin receptor antibodies in Graves’ disease: thyroid peroxidase antibody activity reverts to pretreatment level after carbimazole withdrawal. *Thyroid* 2006;16:1041-5.

# ATA invites You to Join Us at the...



The poster for the 81st Annual Meeting of the American Thyroid Association (ATA) features a green and white color scheme. At the top, it reads "81<sup>ST</sup> Annual Meeting" in a large, bold, black font. Below this is a collage of images: a modern building, a white circular logo with a green thyroid gland shape, a scenic view of a resort, and a dining table. The text "AMERICAN THYROID ASSOCIATION" is centered in a serif font, with "FOUNDED 1923" underneath. The dates "OCTOBER 26-30, 2011" are prominently displayed in a large, bold, black font. At the bottom, the location "Renaissance Esmeralda Resort and Spa, Indian Wells, California" and the website "www.thyroid.org" are listed.

The American Thyroid Association is the leading organization focused on thyroid biology and the prevention and treatment of thyroid disorders through excellence and innovation in research, clinical care, education, and public health.

At the 81<sup>st</sup> Annual Meeting of the American Thyroid Association (ATA), attendees will experience top-notch educational sessions, great networking opportunities and unmatched collegiality -- all under one-roof.

Nestled at the base of the majestic Santa Rosa Mountains in Indian Wells near Palm Springs, CA, the Renaissance Esmeralda Resort & Spa is the Sonoran Desert's finest oasis, a perfect setting for ATA attendees from around the world to meet.

Chaired by Drs. Anthony Hollenberg and Martha Zeiger, the ATA Program Committee promises to

offer the outstanding agenda expected by those who choose the ATA meeting as their 'favorite' scientific educational experience - year after year. Past attendees attest to the unmatched excellence and environment of the ATA meeting noting:

- "Great combination of clinical and basic research"
- "Presentations and posters are excellent"
- "Well organized and top notch"
- "The science improves every year"

**WHY SHOULD YOU ATTEND?** Earn CME credits, hear innovative talks on clinical topics, participate in interactive sessions, develop professionally with state of the art information, and meet with friends and colleagues.

**WHO WILL BE THERE?** The community of endocrinologists, internists, surgeons, basic scientists, nuclear medicine scientists, pathologists, endocrine fellows and nurses, physician assistants and other health care professionals who wish to broaden and update their knowledge of the thyroid gland and its disorders. Clinical, Basic and Surgical Fellows will have a customized educational track to enhance their meeting experience.

## REGISTRATION

ATA meeting registration is open to all health care professionals interested in broadening their knowledge of the thyroid gland and its disorders. **Visit the ATA website for registration details and meeting information as available at [www.thyroid.org](http://www.thyroid.org).**

## HOTEL

Book your hotel reservation now and mention the ATA to receive the special group rate. Renaissance Esmeralda Resort & Spa, 44-400 Indian Wells Lane, Indian Wells, CA 92210; 760-773-4444 or 800-446-9875.

## American Thyroid Association Short Call for Abstracts



AMERICAN  
THYROID  
ASSOCIATION  
FOUNDED 1923

[www.thyroid.org](http://www.thyroid.org)

### Short Call Abstract Submission Deadlines

- Short call Site opens – Wednesday, September 7, 2011
- Site closes – Wednesday, September 21, 2011
- Notification of acceptance: September 30, 2011
  - Online confirmation by corresponding author is required.

### ATA Abstract Submission Policy and Responsibilities of the Author:

The ATA requests submission of abstracts considered for ATA scientific meetings to feature new data to be presented as posters or oral presentations. The ATA goal is to provide the audience and the media with new data that are unpublished (in print or electronic) which are being publicly presented for the first time. Authors are asked to strictly comply with this requirement; data that are to become available to the public *in the setting of a national or international meeting* before their presentation at the ATA meeting are not eligible for presentation at the ATA meeting. Data may be submitted for publication before or after abstract submission to the ATA. However, data accepted for publication prior to the ATA meeting would REQUIRE the authors to request the publisher to embargo their publication (electronic and print) until **8:00 am local time the first day of the meeting**, or would REQUIRE the authors to withdraw their abstract from the ATA meeting. Many editors are favorable to embargo requests because of the attention that may be drawn to the publication after original presentation of the data at a major meeting. Further, the authors are welcome to announce the date and place of their anticipated publication if known. Authors that do not comply with this policy may be restricted from future abstract submissions for a term to be determined by the ATA Executive Committee. Arbitration, if needed, will occur via the ATA Board of Directors. **Abstracts are reviewed in confidence by the ATA program committee with possible ad hoc members.**

### Additional policies:

- **CHARACTER LIMIT:** There is a limit of 2,245 characters (approx. 300 words) for the text of your submission.
- Authors of accepted posters are required to be present during the assigned poster sessions.
- Scientific materials presented at the ATA Annual Meeting must not have been submitted for publication at the time of abstract submission or presented at a scientific meeting before the 81<sup>st</sup> Annual Meeting of the ATA (local and regional meetings excluded).
- All abstracts must be filed electronically via the American Thyroid Association website ([www.thyroid.org](http://www.thyroid.org)). Submissions will not be accepted by fax or mail.
- All materials must arrive on or before the abstract deadlines noted above.
- Authorship on multiple abstracts is permitted.

### Short Call Abstracts

- Short Call Abstracts are reserved for the presentation of the very latest, important thyroid-related research with high impact. Submission of a Short Call Abstract does not guarantee acceptance for presentation.
- Only Six (6) Short Call Abstracts will be selected for 10-minute oral presentations during a special symposium. Selected additional Short Call Abstracts may be presented as special posters. All other submissions will not be published.
- Acceptance notices for those selected will be e-mailed on or before September 30, 2011. Online confirmation is required.

## American Thyroid Association

*Dedicated to Scientific Inquiry, Clinical Excellence, Public Service, Education, and Collaboration*

*ATA Online Seminars: Education When and Where You Want It*

**REGISTER NOW FOR THE SEPTEMBER 20 WEBINAR AT [WWW.THYROID.ORG](http://WWW.THYROID.ORG)**

**LIVE COURSES FREE TO FELLOWS**

### **New Technologies and Techniques in Thyroid Surgery**

**Tuesday, September 20, 2011; 11:00 AM ET**

#### **William B. Inabnet, III, MD, FACS**

Mount Sinai Medical Center

1 CME Credit Available

The field of thyroid surgery has experienced numerous advances since the advent of laparoscopic surgery in the early 1990's. This webinar will provide a comprehensive update on new techniques and technology in thyroid surgery with an emphasis on minimally invasive and video-endoscopic approaches to the thyroid gland. A pathway for safely introducing new techniques to the clinical realm will be discussed. [Learn More...](#)

**Costs:** \$119 for ATA members per webinar/\$149 for non-members per webinar. **Free registration for all fellows who sign up for the 9/20/2011, 11:00 AM ET live webinar.**

Fellows should contact the ATA at [thyroid@thyroid.org](mailto:thyroid@thyroid.org) to receive the complimentary registration code to participate.



**Target Audience (Who Should Attend):** ATA webinars are designed for endocrinologists, internists, surgeons, basic scientists, nuclear medicine scientists, pathologists, endocrine and surgery fellows, nurses, physician assistants and other health care professionals who wish to broaden and update their knowledge of the thyroid gland and its disorders including clinical management guidelines and recent advances in thyroidology.

**Learning Objectives:** At the conclusion of ATA webinars, attendees should be able to:

- Describe state-of-the art findings on the mechanisms, prevention, diagnosis, and management of thyroid disorders and cancer
- Explain the latest clinical management guidelines to benefit patient care and the expertise of the clinician in practice
- Describe the impact of health policy, environmental factors, genetic factors, and non-thyroidal conditions on thyroid disorders and cancer
- Explain new treatment options for thyroid disorders and cancer in patient care
- Identify opportunities for increasing education and collaboration to further understand thyroid disorders, thyroid cancer and managing patient care

**Disclosures:** Disclosures for presenting faculty and content controllers will be provided to attendees verbally or on-screen during the live webinar activity.

**Accreditation and Designation Statement:** intellyst® Medical Education is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians. intellyst® Medical Education designates each of these live activities for a maximum of 1 AMA PRA Category 1 Credit™. Physicians should only claim credit commensurate with the extent of their participation in the activity.

**Recorded webinars available for webinars by Drs David Cooper, Michael McDermott, Steven Sherman, Ralph Tufano and William Inabnet at [www.thyroid.org](http://www.thyroid.org)**

**REGISTER FOR THE ATA ANNUAL MEETING NOW**

**Don't miss out on the discounted ground transportation shuttle offer. Visit [www.thyroid.org](http://www.thyroid.org)**



**81<sup>ST</sup> Annual Meeting**  
Renaissance Esmeralda Resort and Spa  
Indian Wells, California • [www.thyroid.org](http://www.thyroid.org)  
**OCTOBER 26-30, 2011**

  
AMERICAN  
THYROID  
ASSOCIATION  
FOUNDED 1923