

Clinical Activity Score

- Add 1 point for each finding
- Symptoms
 - Pain or pressure in a periorbital or retroorbital distribution
 - Pain with upward, downward, or lateral eye movement
- Signs
 - Swelling of the eyelids
 - Redness of the eyelids
 - Conjunctival injection
 - Chemosis
 - Inflammation of the caruncle or plica
- Changes
 - Increase in measured proptosis \geq 2 mm over 1-3 months
 - Decrease in eye movement limit of $\geq 8^\circ$ over 1-3 months
 - Decrease in visual acuity (2 Snellen chart lines) over 1-3 months

Clinical Activity Score

- Score ≥ 4
 - Positive predictive value 80%
 - Negative predictive value 64%

Basic NOSPECS Classification

- N = No symptoms or signs
- O = Only signs
- S = Soft tissue involvement
- P = Proptosis
- E = Extraocular muscle involvement
- C = Corneal involvement
- S = Sight loss due to optic nerve compression

Modified NOSPECS Classification

- Class 0 = No symptoms or signs
- Class I = Only signs
- Class II = Soft tissue involvement
 - o. Absent
 - a. Minimal
 - b. Moderate
 - c. Marked
- Class III = Proptosis
 - o. Absent
 - a. Minimal
 - b. Moderate
 - c. Marked

Modified NOSPECS Classification

- Class IV = Extraocular muscle involvement
 - o. Absent
 - a. Limitation of motion in extremes of gaze
 - b. Evident restriction of motion
 - c. Fixation of globe or globes
- Class V = Corneal involvement
 - o. Absent
 - a. Stippling of the cornea
 - b. Ulceration
 - c. Clouding, necrosis, perforation

Modified NOSPECS Classification

- Class VI = Sight loss due to optic nerve compression
 - o. Absent
 - a. Visual acuity 0.63-0.5
 - b. Visual acuity 0.4-0.1
 - c. Visual acuity < 0.1 - no light perception